

Quantum Lax scheme for Ruijsenaars models

Branislav Jurčo^{*1} and Peter Schupp^{**}

^{*}Max-Planck-Institut für Mathematik

Vivatgasse 7

D-53111 Bonn, Germany

^{**}Sektion Physik

Universität München

Theresienstr. 37

D-80333 München, Germany

Abstract

We develop a quantum Lax scheme for IRF models and difference versions of Calogero-Moser-Sutherland models introduced by Ruijsenaars. The construction is in the spirit of the Adler-Kostant-Symes method generalized to the case of face Hopf algebras and elliptic quantum groups with dynamical R-matrices.

1 Introduction

The Hamiltonian of the (relativistic) Ruijsenaars model [4] for two particles on a line with coordinates x_1 and x_2 acts on a wave function in a manifestly non-local way as

$$\mathcal{H}\psi(\lambda) = \frac{\theta(\frac{c\eta}{2} - \lambda)}{\theta(-\lambda)}\psi(\lambda - \eta) + \frac{\theta(\frac{c\eta}{2} + \lambda)}{\theta(\lambda)}\psi(\lambda + \eta), \quad (1)$$

where $\lambda = x_1 - x_2$ is the relative coordinate, $c \in \mathbb{C}$ is a coupling constant, η is the relativistic deformation parameter, and θ is the Jacobi theta function. The Hamiltonian apparently contains shift operators t_i that generate a one-dimensional graph:

¹e-mail: jurco@mpim-bonn.mpg.de



Relative to a fixed vertex λ the vertices of this (ordered) graph are at points $\eta \cdot \mathbb{Z} \in \mathbb{R}$. This picture generalizes arbitrary ordered graphs, which we shall consider in the following. We will continue to fix a vertex λ to avoid a continuous family of disconnected graphs.

1.1 Face algebras

The Hilbert space of the model are vector spaces on paths of fixed length on the ordered graph. Its operators are elements of a Face Algebra F [6]. This is a novel mathematical structure that naturally incorporates the complicated shifts of the Ruijsenaars model. There are two commuting projection operators $e^i, e_i \in F$ onto bra's and ket's corresponding to each vertex i : $e_i e_j = \delta_{ij} e_i$, $e^i e^j = \delta_{ij} e^i$, $\sum e_i = \sum e^i = 1$. F has a coalgebra structure such that $e_j^i \equiv e^i e_j = e_j e^i$ is a corepresentation:


$$\Delta(e_j^i) = \sum_k e_k^i \otimes e_j^k, \quad \epsilon(e_j^i) = \delta_{ij}; \quad \Rightarrow \quad \Delta(1) = \sum_k e_k \otimes e^k \neq 1 \otimes 1. \quad (2)$$

The latter is a key feature of face algebras and weak C^* -Hopf algebras. The matrix indices of e_j^i are vertices, *i.e.* paths of length zero. In the given setting it is natural to also allow paths of fixed length on a finite oriented graph $\mathcal{G}$ as matrix indices. We shall use capital letters to label paths. A path P has an origin (source) $\cdot P$, an end (range) $P \cdot$ and a length $\#P$. Two paths Q, P can be concatenated to form a new path $Q \cdot P$, if the end of the first path coincides with the start of the second path, *i.e.* if $Q \cdot = \cdot P$. The symbols T_B^A , where $\#A = \#B = \#A' \geq 0$, with relations

$$\Delta(T_B^A) = \sum_{A'} T_{A'}^A \otimes T_B^{A'}, \quad \epsilon(T_B^A) = \delta_{AB}, \quad T_B^A T_D^C = \delta_{A \cdot, C} \delta_{B \cdot, D} T_{B \cdot D}^{A \cdot C} \quad (3)$$

span an object that obeys the axioms of a face algebra. Relations (3) make T_B^A a corepresentation; and the last expression is the rule for combining representations. The axioms of a face algebra can be found in [6].

Pictorial representation:



The dashed paths indicate the F -space(s). Inner paths are summed over.

It is convenient to work with an abstract, universal T , which is the canonical element T_{12} of $U \otimes F$, where U is the dual of F via the pairing $\langle , \rangle$.²

$$\langle T, f \rangle = f, \quad \langle T_1 \otimes T_1, f \rangle = \Delta f, \quad \langle T_1 T_2, f \otimes g \rangle = fg; \quad f, g \in F$$

A face *Hopf* algebra has an antipode S which is denoted by a tilde in the universal tensor formalism: $\langle \tilde{T}, f \rangle = S(f)$. An ordinary Hopf algebra is a special case of a Face Hopf algebra with a single vertex. Ordinary matrix indices correspond to closed loops in that case.

1.2 Boltzmann weights

The axioms [6] for a quasitriangular face algebra are similar to those of a quasitriangular Hopf algebra; there is a universal $R \in U \otimes U$ that controls the non-cocommutativity of the coproduct in U and the non-commutativity of the product in F ,

$$RT_1 T_2 = T_2 T_1 R, \quad \tilde{R} T_2 T_1 = T_1 T_2 \tilde{R}, \quad \tilde{R} \equiv (S \otimes \text{id})(R), \quad (4)$$

however the antipode of R is no longer inverse of R but rather $\tilde{R}R = \Delta(1)$. The numerical “ R -matrix” obtained by contracting R with two face corepresentations is given by the face Boltzmann weight W :

$$\langle R, T_B^A \otimes T_D^C \rangle = R_{BD}^{AC} \equiv W \left(C \begin{matrix} B \\ A \end{matrix} D \right) \sim \begin{array}{c} \text{---} B \text{---} \\ \uparrow \quad \downarrow \\ C \quad \quad D \\ \downarrow \quad \uparrow \\ \text{---} A \text{---} \end{array}$$

The Boltzmann weight is zero unless $C \cdot A$ and $B \cdot D$ are valid paths with common source and range. For $f \in F$ there are two algebra homomorphisms $F \rightarrow U$:

$$R^+(f) = \langle R, f \otimes \text{id} \rangle, \quad R^-(f) = \langle \tilde{R}, \text{id} \otimes f \rangle. \quad (5)$$

R satisfies the YangBaxter Equation $R_{12}R_{13}R_{23} = R_{23}R_{13}R_{12}$. Contracted with $T_B^A \otimes T_D^C \otimes T_F^E$ this expression yields a numerical Yang-Baxter equation with the following pictorial representation [2]:

The inner edges are paths that are summed over.

²Here and in the following we shall frequently suppress the the second index of T ; it corresponds to the F -space. The displayed expressions are short for $\langle T_{12}, f \otimes \text{id} \rangle = f \in F$, $\langle T_{12}T_{13}, f \otimes \text{id} \otimes \text{id} \rangle = \Delta f \in F \otimes F$ and $\langle T_{13}T_{23}, f \otimes g \otimes \text{id} \rangle = fg \in F$.

2 Quantum factorization

Cocommutative functions, like the trace of the T-matrix, provide mutually commutative operators including the Hamiltonian. The following theorem gives for the case of face Hopf algebras what has become known as the “Main theorem” for the solution by factorization of the equations of motion [7, 8, 9, 10]:

Theorem 2.1 (Main theorem for face algebras)

- (i) The set of cocommutative functions, I , is a commutative subalgebra of F .
- (ii) The Heisenberg equations of motion defined by a Hamiltonian $\mathcal{H} \in I$ are of Lax form $i \frac{dT}{dt} = [M^\pm, T]$, with $M^\pm = 1 \otimes \mathcal{H} - m_\pm \in U_\pm \otimes F$, $m_\pm = R^\pm(\mathcal{H}_{(2)}) \otimes \mathcal{H}_{(1)}$; see (5).
- (iii) Let $g_\pm(t) \in U_\pm \otimes F$ be the solutions to the factorization problem $g_-^{-1}(t)g_+(t) = \exp(it(m_+ - m_-)) \in U \otimes F$, then $T(t) = g_\pm(t)T(0)g_\pm(t)^{-1}$ solves the Lax equation; $g_\pm(t)$ are given by $g_\pm(t) = \exp(-it(1 \otimes h)) \exp(it(1 \otimes h - M^\pm(0)))$ and are the solutions to the differential equation $i \frac{d}{dt} g_\pm(t) = M^\pm(t)g_\pm(t)$, $g_\pm(0) = 1$.

3 Dynamical operators

Like we mentioned in the introduction we are interested in the action of the Hamiltonian with respect to a fixed vertex. We fix a vertex with the help of $e^\lambda, e_\lambda \in F$. R-matrices with a fixed vertex are *Dynamical R-matrices*: $R_{12}(\lambda) \equiv \langle R, T_1(\lambda) \otimes T_2 \rangle$, where $T(\lambda)_R^A$ is zero unless the range (end) of path A is equal to the fixed vertex λ :

$$T(\lambda)_B^A = T_B^A e^\lambda \quad \sim \quad \begin{array}{c} \xrightarrow{A} \times^\lambda \\ \uparrow \hspace{1.5cm} \uparrow \\ \text{---} \hspace{0.5cm} \text{---} \\ \downarrow \hspace{1.5cm} \downarrow \\ \xrightarrow{B} \end{array} \quad (6)$$

The formalism naturally includes shifts, as can e.g. be seen in the *Dynamical RTT-equation* [1]

$$R_{12}(\lambda)T_1(\lambda - h_2)T_2(\lambda) = T_2(\lambda - h_1)T_1(\lambda)R_{12}(\lambda - h_3). \quad (7)$$

and the *Dynamical Yang-Baxter equation* [3]

$$R_{12}(\lambda)R_{13}(\lambda - h_2)R_{23}(\lambda) = R_{23}(\lambda - h_1)R_{13}(\lambda)R_{12}(\lambda - h_3). \quad (8)$$

The shift operators h_1, h_2, h_3 can be expressed in terms of e^μ, e_η and their duals E^μ, E_η , if we assume a (local) embedding of the vertices of the graph in $\mathbb{C}^\times$.

The Hamiltonian of the Ruijsenaars model is the trace of a T -matrix, in an appropriate representation. It can be written as a sum of operators that act in subspaces

corresponding to paths ending in a vertex λ : $\mathcal{H} = \sum_{\lambda} \mathcal{H}(\lambda)$ with $\mathcal{H}(\lambda) = \mathcal{H}e^{\lambda} = \sum T(\lambda)_Q^Q$. The pictorial representation of the Hamiltonian is two closed dashed paths (F -space) connected by paths Q of fixed length that are summed over. In $\mathcal{H}(\lambda)$ the end of path Q is fixed. When we look at a representation on Hilbert space the paths Q with endpoint λ that appear in the component $\mathcal{H}(\lambda)$ of the Hamiltonian $\mathcal{H}$ will shift the argument of a state $\psi(\lambda)$ corresponding to the vertex λ to a new vertex corresponding to the starting point of the path Q , exactly as in (1).

Given the Boltzmann weights [5, 11], the Main Theorem provides Lax operators and Lax equations for the Ruijsenaars model [12].

Acknowledgments

We would like to thank Pavel Winternitz for interest and support, and Koji Hasegawa and Jan Felipe van Diejen for interesting discussions. B. J. would like to thank the Alexander von Humboldt Foundation for support.

References

- [1] Felder, G.: Elliptic quantum groups. In Proc. XIth International Congress of Mathematical Physics, D. Iagolnitzer (ed.), Boston: International Press, 1995
- [2] Baxter, R.J.: Exactly solved models in statistical physics. New York: Academic Press, 1982
- [3] Gervais, J.L., Neveu, A., Nucl. Phys. **B 238**, 125 (1984)
- [4] Ruijsenaars, S.N.M., Comm. Math. Phys. **110**, 191 (1987)
- [5] Hasegawa, K., Comm. Math. Phys. **187**, 289 (1997)
- [6] Hayashi, T., Publ. RIMS, Kyoto University **32**, 351 (1996)
- [7] Adler, M., Inv. Math. **50**, 219 (1979); Kostant, B., Inv. Math. **159**, 13 (1980)
- [8] Reyman, A. G., Semenov-Tian-Shansky, M. A.: Group theoretical methods in the theory of finite-dimensional integrable systems. In Encyclopedia of mathematical sciences, Dynamical systems VII: Springer 1993
- [9] Jurčo, B., Schlieker, M., Comm. Math. Phys **185**, 397 (1997)
- [10] Jurčo, B., Schupp, P., Int. J. Mod. Phys. **A 12**, 5735 (1997)
- [11] Jimbo, M., Miwa, T., Okado, M., Nucl. Phys. **B 300**, 74 (1988)
- [12] Jurčo, B., Schupp, P., J. Math. Phys. **39**, 3577 (1998)